arXiv:hep-ex/0604052v1 27 Apr 2006

Pion Form Factor at SND (*new edition*).

M.N. Achasov [a*], K.I.Beloborodov[ab], A.V.Berdyugin[a], A.G.Bogdanchikov[a], A.V.Bozhenok[ab],
A.D.Bukin[a], D.A.Bukin[a], T.V.Dimova[a], V.P.Druzhinin[ab], V.B.Golubev[ab], A.A.Korol[a], S.V.Koshuba[a],
E.V.Pakhtusova[a], S.I.Serednyakov[ab], Yu.M.Shatunov[ab], V.A.Sidorov[a], Z.K.Silagadze[ab], A.N.Skrinsky[a],
Yu.A.Tikhonov[ab], and A.V.Vasiljev[ab],

[a]Budker Institute of Nuclear Physics, Siberian Branch of the Russian Academy of Sciences,
11 Lavrentyev,Novosibirsk, 630090, Russia

[b]Novosibirsk State University, 630090, Novosibirsk, Russia

The update of the $e^+e^- \to \pi^+\pi^-$ process cross section, measured in the energy region $\sqrt{s} < 1$ GeV with SND detector at VEPP-2M collider is presented.

The $e^+e^- \to \pi^+\pi^-$ process cross section was measured by SND [2] detector at VEPP-2M [1] collider in the energy region below 1 GeV [3]. The measurement was based on about 5×10^6 $e^+e^- \to \pi^+\pi^-$ events and has systematic accuracy 1.3%.

Calculation of the $e^+e^- \to e^+e^-\gamma$, $\pi^+\pi^-\gamma$ and $\mu^+\mu^-\gamma$ processes cross sections plays quite an important role in form factor measurement. It is necessary for the luminosity measurement ($e^+e^- \to e^+e^-\gamma$), for background subtraction ($e^+e^- \to \mu^+\mu^-\gamma$), for radiative corrections and detection efficiency determination ($e^+e^- \to \pi^+\pi^-\gamma$). Recently it was found, that $e^+e^- \to \pi^+\pi^-\gamma$ and $e^+e^- \to \mu^+\mu^-\gamma$ Monte Carlo events generators based on Ref.[5,4] were not quite correct. So we performed reanalysis of SND data using MCGPJ generator [6], which is based on the same articles. As a result of reanalysis the corrections to the measured $e^+e^- \to \pi^+\pi^-$ cross section [3] were obtained (fig.1). After the corrections the cross section values decreased by two systematic errors.



Figure 1. Corrections δ to the $e^+e^- \to \pi^+\pi^-$ cross section. • and ∘ are corrections due to inaccuracy of the $e^+e^- \to \mu^+\mu^-\gamma$ and $\pi^+\pi^-\gamma$ events generator, $\star$ is the total correction.

The comparison of corrected SND result with the CMD-2 precision measurements [7,8] is shown in fig.2 and 3. In the energy region $\sqrt{s}$ from 600 to 1000 MeV the average deviation

$$\Delta = (1 - \langle\sigma_{CMD-2}/\sigma_{SND}\rangle) \times 100\% \qquad (1)$$

between the SND (σ_{SND}) and CMD-2 (σ_{CMD-2}) measurements is equal to $-1.4 \pm 1.5\%$ for the published [7] CMD-2 data and $-0.3 \pm 1.6\%$ for the new, not yet published but already reported, CMD-2 result [8]. For uncorrected SND cross section Δ was equal to $1.4 \pm 1.5\%$ [3] and $2.5 \pm 1.6\%$ [9] respectively. The errors include both system-

---

[*] E-mail:achasov@inp.nsk.su.



Figure 2. The ratio of the $e^+e^- \to \pi^+\pi^-$ cross section obtained in CMD-2 [7] and SND (this talk) experiments. The shaded area shows the joint systematic error.



Figure 3. The ratio of the $e^+e^- \to \pi^+\pi^-$ cross section obtained in CMD-2 [8] and SND (this talk) experiments. The shaded area shows the joint systematic error.

atic and statistic uncertainties of the CMD-2 and SND measurements.

The table of corrected cross sections and results of theoretical fit to the data will be published soon.

The authors are grateful to G.V. Fedotovich, F.V. Ignatov, A.L. Sibidanov for useful discussions. The work is supported in part by grants Sci.School-1335.2003.2, RFBR 04-02-16181-a, 04-02-16184-a, 05-02-16250-a, 06-02-16192-a.

REFERENCES

1. A.N. Skrinsky, in Proceedings of Workshop on physics and detectors for DAΦNE, Frascati, Italy, April 4-7, 1995, p.3.
2. M.N. Achasov et al., Nucl. Instr. and Meth. A 449 (2000) 125
3. M.N. Achasov et al., Zh. Eksp. Teor. Fiz. B 128 (2005) 1201; JETP 101 (2005) 1053.
4. A.B. Arbuzov et al., JHEP 10 (1997) 001
5. A.B. Arbuzov et al., JHEP 10 (1997) 006
6. A.B. Arbuzov et al., Report of Budker INP 2004-70, Novosibirsk (2004); hep-ph/0504233; Computer code was got from `http://cmd.inp.nsk.su/~sibid/radnsk.tar.bz2`
7. R.R. Akhmetshin et al, Phys. Lett. B 527 (2002) 161; Phys. Lett. B 578 (2004) 285
8. I.B. Logashenko et al., in Proceedings of International European Conference on High Energy Physics, Lisboa, Portugal, July 21-27, 2005; A.L. Sibidanov et al., in Proceedings of 11 International Conference on Hadron Spectroscopy, Rio-De-Janeiro, Brazil, August 21-26, 2005; I.B. Logashenko, this Proceedings
9. M.N. Achasov, in Proceedings of Conf. "Physics of fundamental interactions", dedicated to 60th anniversary of ITEP (Session of RAS Nuclear Physics Department 2005), Moscow, December 5-9, 2005